

Existence of the σ -Meson below 1 GeV and the $f_0(1500)$ Glueball

Yu.S. Surovtsev^{a,1}, D. Krupa^b, M. Nagy^b

^a*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research,
Dubna 141 980, Moscow Region, Russia,*

^b*Institute of Physics, Slov.Acad.Sci., Dúbravská cesta 9, 842 28 Bratislava, Slovakia*

Abstract

On the basis of a simultaneous description of the isoscalar s -wave channel of the $\pi\pi$ scattering (from the threshold up to 1.9 GeV) and of the $\pi\pi \rightarrow K\bar{K}$ process (from the threshold to ~ 1.4 GeV) in the model-independent approach, a confirmation of the σ -meson at ~ 665 MeV and an indication for the glueball nature of the $f_0(1500)$ state are obtained. It is shown that the large $\pi\pi$ -background, usually obtained, combines, in reality, the influence of the left-hand branch-point and the contribution of a very wide resonance at ~ 665 MeV. The coupling constants of the observed states with the $\pi\pi$ and $K\bar{K}$ systems and lengths of the $\pi\pi$ and $K\bar{K}$ scattering are obtained.

PACS: 14.40.Cs, 11.80.Gw, 12.39.Pn, 13.75.Lb

Keywords: Multichannel states; Riemann surface; Pole clusters

¹E-mail address: surovcev@thsun1.jinr.ru

1. A problem of scalar mesons is most troublesome and long-lived in the light meson spectroscopy. Among difficulties in their understanding there is the one related to a strong model-dependence of information on multichannel states obtained in analyses based on the specific dynamic models or using an insufficiently-flexible representation of states (*e.g.*, the standard Breit – Wigner form). Earlier, we have shown [1] that an inadequate description of multichannel states gives not only their distorted parameters when analyzing data but also can cause the fictitious states when one neglects important (even energetic-closed) channels. In this paper we are going, conversely, to demonstrate that the large background (*e.g.*, that happens in analyzing $\pi\pi$ scattering), can hide low-lying states, even such important for theory as a σ -meson [2]. The latter is required by most of the models (like the linear σ -models or the Nambu – Jona-Lasinio models [3]-[6]) for spontaneous breaking of chiral symmetry. Recent new analyses of the old and new experimental data found a possible candidate for that state [7]-[11]. However, these analyses use either the Breit – Wigner form (even if modified) or specific forms of interactions; therefore, there one cannot talk about a model independence of results. Besides, in these analyses, a large $\pi\pi$ -background is obtained.

A model-independent information on multichannel states can be obtained on the basis of the first principles (analyticity, unitarity) immediately applied to analyzing experimental data. The way of realization is a consistent allowance for the nearest singularities on all sheets of the Riemann surface of the S -matrix. Earlier, we have proposed this method for 2- and 3-channel resonances and developed the concept of standard clusters (poles on the Riemann surface) as a qualitative characteristic of a state and a sufficient condition of its existence [1, 12]. The cluster kind is related to the state nature. At all events, we can, in a model-independent manner, discriminate between bound states of particles and the ones of quarks and gluons [1, 13], qualitatively predetermine the relative strength of coupling of a state with the considered channels, and obtain an indication on its glunium nature.

2. Here we restrict ourselves to a 2-channel simultaneous consideration of the coupled processes $\pi\pi \rightarrow \pi\pi, K\bar{K}$. Therefore, we have the 2-channel S -matrix determined on the 4-sheeted Riemann surface. The matrix elements $S_{\alpha\beta}$, where $\alpha, \beta = 1(\pi\pi), 2(K\bar{K})$, have the right-hand cuts along the real axis of the s -plane, starting at $4m_\pi^2$ and $4m_K^2$, and the left-hand cuts, beginning at $s = 0$ for S_{11} and at $4(m_K^2 - m_\pi^2)$ for S_{22} and S_{12} . We number the Riemann-surface sheets according to the signs of analytic continuations of the channel momenta $k_1 = (s/4 - m_\pi^2)^{1/2}$, $k_2 = (s/4 - m_K^2)^{1/2}$ as follows: signs $(\text{Im}k_1, \text{Im}k_2) = ++, +-, --, +-$ correspond to the sheets I, II, III, IV.

To elucidate the resonance representation on the Riemann surface, we express analytic

continuations of the matrix elements to the unphysical sheets $S_{\alpha\beta}^L$ ($L = II, III, IV$) in terms of them on the physical sheet $S_{\alpha\beta}^I$. The latter have, except for the real axis, only zeros corresponding to resonances. Using the reality property of the analytic functions and the 2-channel unitarity, one can obtain

$$\begin{aligned} S_{11}^{II} &= \frac{1}{S_{11}^I}, & S_{11}^{III} &= \frac{S_{22}^I}{\det S^I}, & S_{11}^{IV} &= \frac{\det S^I}{S_{22}^I}, \\ S_{22}^{II} &= \frac{\det S^I}{S_{11}^I}, & S_{22}^{III} &= \frac{S_{11}^I}{\det S^I}, & S_{22}^{IV} &= \frac{1}{S_{22}^I}, \\ S_{12}^{II} &= \frac{iS_{12}^I}{S_{11}^I}, & S_{12}^{III} &= \frac{-S_{12}^I}{\det S^I}, & S_{12}^{IV} &= \frac{iS_{12}^I}{S_{22}^I}, \end{aligned} \quad (1)$$

Here $\det S^I = S_{11}^I S_{22}^I - (S_{12}^I)^2$. These formulae immediately give the resonance representation by poles and zeros on the 4-sheeted Riemann surface. One must discriminate between three types of 2-channel resonances described by a pair of conjugate zeros on sheet I: **(a)** in S_{11} , **(b)** in S_{22} , **(c)** in each of S_{11} and S_{22} . As is seen from (1), to the resonances of types **(a)** and **(b)** one has to make correspond a pair of complex conjugate poles on sheet III, shifted relative to a pair of poles on sheet II and IV, respectively. To the states of type **(c)** one must make correspond two pairs of conjugate poles on sheet III. A resonance of every type is represented by a pair of complex-conjugate clusters (of poles and zeros on the Riemann surface) of size typical of strong interactions. The cluster kind is related to the state nature. The resonance, coupled strongly with the $\pi\pi$ channel, is described by the cluster of type **(a)**; the resonance, coupled strongly with the $K\bar{K}$ and weakly with $\pi\pi$ channel (say, an $s\bar{s}$ state), by the cluster of type **(b)**; the flavour singlet (*e.g.* glueball) must be represented by the cluster of type **(c)** as a necessary condition.

For the simultaneous analysis of experimental data on the coupled processes it is convenient to use the Le Couteur-Newton relations [14, 15] expressing the S -matrix elements of all coupled processes in terms of the Jost matrix determinant $d(k_1, k_2)$, the real analytic function with the only square-root branch-points at $k_i = 0$. Earlier, this was done by us in the 2-channel consideration [12] with the uniformizing variable $z = (k_1 + k_2)/\sqrt{m_K^2 - m_\pi^2}$ which was proposed in Ref.[16] and maps the 4-sheeted Riemann surface with two unitary cuts, starting at $4m_\pi^2$ and $4m_K^2$, onto the plane. Note that other authors have been also applied the parametrizations with using the Jost functions at analyzing the s -wave $\pi\pi$ scattering in the one-channel approach [17] and in the 2-channel one [13]. In latter work, the uniformizing variable k_2 has been used, therefore, their approach cannot be employed near by the $\pi\pi$ threshold.

When analyzing the processes $\pi\pi \rightarrow \pi\pi, K\bar{K}$ by the above methods in the 2-channel approach, two states ($f_0(975)$ and $f_0(1500)$) were found to be sufficient ($\chi^2/\text{ndf} \approx 1.00$). However, there the large $\pi\pi$ -background has been obtained. A character of the repre-

sensation of the background (the pole of 2nd order on the real axis on sheet II and the corresponding zero on sheet I) suggests that a wide light state is possible to be hidden in the background. To check this, one must work out the background in some detail.

Now we will take, in the uniformizing variable, into account also the left-hand branch-point at $s = 0$. We use the uniformizing variable

$$v = \frac{m_K \sqrt{s - 4m_\pi^2} + m_\pi \sqrt{s - 4m_K^2}}{\sqrt{s(m_K^2 - m_\pi^2)}}, \quad (2)$$

which maps the 4-sheeted Riemann surface, having (in addition to two above-indicated unitary cuts) also the left-hand cut starting at $s = 0$, onto the v -plane. In Fig.1, the plane of the uniformizing variable v for the $\pi\pi$ -scattering amplitude is depicted. The Roman numerals (I, ..., IV) denote the images of the corresponding sheets; the thick line represents the physical region; the points i , 1 and $b = \sqrt{(m_K + m_\pi)/(m_K - m_\pi)}$ correspond to the $\pi\pi$, $K\bar{K}$ thresholds and $s = \infty$, respectively; the shaded intervals $(-\infty, -b]$, $[-b^{-1}, b^{-1}]$, $[b, \infty)$ are the images of the corresponding edges of the left-hand cut. The depicted positions of poles (*) and of zeros (o) give the representation of the type (a) resonance in S_{11} .

On v -plane the Le Couteur-Newton relations are [12, 16]

$$S_{11} = \frac{d(-v^{-1})}{d(v)}, \quad S_{22} = \frac{d(v^{-1})}{d(v)}, \quad S_{11}S_{22} - S_{12}^2 = \frac{d(-v)}{d(v)}. \quad (3)$$

The condition of the real analyticity implies $d(-v^*) = d^*(v)$ for all v , and the unitarity needs the following relations to hold true for the physical v -values: $|d(-v^{-1})| \leq |d(v)|$, $|d(v^{-1})| \leq |d(v)|$, $|d(-v)| = |d(v)|$.

The d -function that on the v -plane already does not possess branch-points is taken as $d = d_B d_{res}$, where $d_B = B_\pi B_K$; B_π contains the possible remaining $\pi\pi$ -background contribution, related to exchanges in crossing channels; B_K is that part of the $K\bar{K}$ background which does not contribute to the $\pi\pi$ -scattering amplitude. The function $d_{res}(v)$ represents the contribution of resonances, described by one of three types of the pole-zero clusters, *i.e.*, except for the point $v = 0$, it consists of zeros of clusters:

$$d_{res} = v^{-M} \prod_{n=1}^M (1 - v_n^* v) (1 + v_n v), \quad (4)$$

where M is the number of pairs of the conjugate zeros.

3. Here we analyze simultaneously the available experimental data on the $\pi\pi$ -scattering [18] and the process $\pi\pi \rightarrow K\bar{K}$ [19] in the channel with $I^G J^{PC} = 0^+ 0^{++}$.

To obtain the satisfactory description of the $\pi\pi$ scattering from the threshold to 1.89 GeV, we have taken $B_\pi = 1$, and three states turned out to be sufficient: the two ones

of the type (a) ($f_0(665)$ and $f_0(980)$) and $f_0(1500)$ of the type (c). The following zero positions on the v -plane, corresponding to these resonances, have been established:

$$\begin{aligned}
\text{for } f_0(665) : \quad & v_1 = 1.36964 + 0.208632i, & v_2 = 0.921962 - 0.25348i, \\
\text{for } f_0(980) : \quad & v_3 = 1.04834 + 0.0478652i, & v_4 = 0.858452 - 0.0925771i, \\
\text{for } f_0(1500) : \quad & v_5 = 1.2587 + 0.0398893i, & v_6 = 1.2323 - 0.0323298i, \\
& v_7 = 0.809818 - 0.019354i, & v_8 = 0.793914 - 0.0266319i.
\end{aligned}$$

Here for the $\pi\pi$ phase shift δ_1 and the elasticity parameter η , 113 and 50 experimental points [18], respectively, are used; when rejecting the points at 0.61, 0.65, and 0.73 GeV for δ_1 and at 0.99, 1.65, and 1.85 GeV for η , which give an anomalously large contribution to χ^2 , we obtain for χ^2/ndf the values 2.7 and 0.72, respectively; the total χ^2/ndf in the case of the $\pi\pi$ scattering is 1.96.

With the presented picture, the satisfactory description for the modulus of the $\pi\pi \rightarrow K\bar{K}$ matrix element $|S_{12}|$ is given from the threshold to ~ 1.4 GeV (Fig.2). Here 35 experimental points [19] are used; $\chi^2/\text{ndf} \approx 1.11$ when eliminating the points at 1.002, 1.265, and 1.287 GeV (with especially large contribution to χ^2). However, for the phase shift $\delta_{12}(s)$, slightly excessive curve is obtained. Therefore, keeping the *parameterless* description of the $\pi\pi$ background, one must take into account the part of the $K\bar{K}$ background that does not contribute to the $\pi\pi$ -scattering amplitude. Note that on the v -plane, S_{11} has no cuts; however, the amplitudes of the processes $K\bar{K} \rightarrow \pi\pi, K\bar{K}$ do have the cuts which arise from the left-hand cut on the s -plane, starting at $s = 4(m_K^2 - m_\pi^2)$. This left-hand cut will be neglected in the Riemann-surface structure, and the contribution on the cut will be taken into account in the $K\bar{K}$ background as a pole on the real s -axis on the physical sheet in the sub- $K\bar{K}$ -threshold region; on the v -plane, this pole gives two poles on the unit circle in the upper half-plane, symmetric to each other with respect to the imaginary axis, and two zeros, symmetric to the poles with respect to the real axis, i.e., one additional parameter is introduced (a position p of the pole on the unit circle). Therefore, for B_K we take the form

$$B_K = v^{-4}(1 - pv)^4(1 + p^*v)^4. \quad (5)$$

Fourth power in (5) is stipulated by the following. First, a pole on the real s -axis on sheet I in S_{22} is accompanied by a pole on sheet II at the same s -value (as it is seen from eqs. (1)); on the v -plane, this implies the pole of second order. Second, for the s -channel process $\pi\pi \rightarrow K\bar{K}$, the crossing u - and t -channels are the $\pi - K$ and $\bar{\pi} - K$ scattering; this results in the additional doubling of the multiplicity of the indicated pole on the v -plane. The expression (5) does not contribute to S_{11} , i.e. the parameterless description of the $\pi\pi$

background is kept. A satisfactory description of the $\delta_{12}(\sqrt{s})$ (Fig.3) is obtained to ~ 1.52 GeV with the parameter $p = 0.948201 + 0.31767i$ (that corresponds to the position of the pole on the s -plane at $s = 0.434\text{GeV}^2$). Here 59 experimental points [19] are considered; $\chi^2/\text{ndf} \approx 3.05$ when eliminating the points at 1.117, 1.247, and 1.27 GeV. The total χ^2/ndf for four analyzed quantities to describe the processes $\pi\pi \rightarrow \pi\pi, K\bar{K}$ is 2.12; the number of adjusted parameters is 17, where they all (except a single relating to the $K\bar{K}$ background) are positions of poles describing resonances.

In Table 1, the obtained poles on the corresponding sheets are cited on the complex energy plane ($\sqrt{s_r} = E_r - i\Gamma_r$). Since, for wide resonances, values of masses and widths are very model-dependent, it is reasonable to report characteristics of pole clusters which must be rather stable for various models.

Now we can calculate the constants of the obtained-state couplings with the $\pi\pi - "1"$ and $K\bar{K} - "2"$ systems through the residues of amplitudes at the pole on sheet II. Expressing the T -matrix via the S -matrix as $S_{ii} = 1 + 2i\rho_i T_{ii}$, $S_{12} = 2i\sqrt{\rho_1\rho_2}T_{12}$, where $\rho_i = \sqrt{(s - 4m_i^2)/s}$, taking the resonance part of the amplitude in the form: $T_{ij}^{res} = \sum_r g_{ir}g_{rj}D_r^{-1}(s)$, where $D_r(s)$ is an inverse propagator ($D_r(s) \propto s - s_r$), and denoting the coupling constants with the $\pi\pi$ and $K\bar{K}$ systems through g_1 and g_2 , respectively, we obtain for $f_0(665)$: $g_1 = 0.7477 \pm 0.095$ GeV and $g_2 = 0.834 \pm 0.1$ GeV, for $f_0(980)$: $g_1 = 0.1615 \pm 0.03$ GeV and $g_2 = 0.438 \pm 0.028$ GeV, for $f_0(1500)$: $g_1 = 0.899 \pm 0.093$ GeV.

Let us indicate also scattering lengths calculated in our approach. For the $K\bar{K}$ scattering, we obtain $a_0^0(K\bar{K}) = -1.188 \pm 0.13 + (0.648 \pm 0.09)i$, $m_{\pi+}^{-1}$. A presence of the imaginary part in $a_0^0(K\bar{K})$ reflects the fact, that already at the threshold of the $K\bar{K}$ scattering, other channels ($2\pi, 4\pi$ etc.) are opened.

For the $\pi\pi$ scattering, we obtain: $a_0^0 = 0.27 \pm 0.06$, $m_{\pi+}^{-1}$. Compare with results of some other works both theoretical and experimental: the value 0.26 ± 0.05 (L. Rosselet et al.[18]), obtained in the analysis of the decay $K \rightarrow \pi\pi e\nu$ with using Roy's model; 0.24 ± 0.09 (A.A. Bel'kov et al.[18] from analysis of the process $\pi^- p \rightarrow \pi^+ \pi^- n$ with using the effective range formula; 0.23 (S. Ishida et al.[9], modified approach to analysis of $\pi\pi$ scattering with using Breit-Wigner forms; 0.16 (S. Weinberg [20], current algebra (non-linear σ -model)); 0.20 (J. Gasser, H. Leutwyler [21], the theory with the non-linear realization of chiral symmetry); 0.26 (M.K. Volkov [22], the theory with the linear realization of chiral symmetry).

We have here presented model-independent results: the pole positions, coupling constants and scattering lengths. The formers can be used further for calculating masses and widths of these states in various models.

If we suppose, that the obtained state $f_0(665)$ is the σ -meson, then from the known relation of the σ -model between the coupling constant of the σ with the $\pi\pi$ -system and masses $g_{\sigma\pi\pi} = (m_\sigma^2 - m_\pi^2)/\sqrt{2}f_{\pi^0}$ (here f_{π^0} is the constant of the weak decay of the π^0 : $f_{\pi^0} = 93.1$ MeV), we obtain $m_\sigma \approx 342$ MeV. That small value of the σ -mass can be a result of the mixing with the $f_0(980)$ state [23].

4. In the present model-independent approach, a satisfactory simultaneous description of the isoscalar s -wave channel of the processes $\pi\pi \rightarrow \pi\pi, K\bar{K}$ from the thresholds to the energy values, where the 2-channel unitarity is valid, is obtained. A parameterless description of the $\pi\pi$ background is first given by allowance for the left-hand branch-point in the proper uniformizing variable. Thus, a model-independent confirmation of the state, already discovered in other works [8]-[11] (or pretending to this discovery) and denoted in the PDG issues by $f_0(400 - 1200)$ [2], is obtained. Three states ($f_0(665)$ – σ -meson, $f_0(980)$ and $f_0(1500)$) are sufficient to describe the analyzed data.

The discovery of the $f_0(665)$ state solves one important mystery of the scalar-meson family that is related to the Higgs boson of the hadronic sector. This is a result of principle, because the schemes of the nonlinear realization of the chiral symmetry have been considered which do without the Higgs mesons. An existence of the low-lying state with the properties of the σ -meson and the obtained value of the $\pi\pi$ -scattering length seem to suggest the linear realization of chiral symmetry. Remember that this simple and beautiful mechanism works also in other fields of physics, *e.g.*, in superconductivity and that the effective Lagrangians obtained on the basis of this mechanism (the Nambu – Jona-Lasinio and other models) describe perfectly the ground states and related phenomena. The only weak link of this approach was the absence of the σ -meson below 1 GeV.

Let us notice that the character of the $f_0(665)$ pole-cluster (namely, a considerable shift of the pole on sheet III towards the imaginary axis) can point to the unconsidered channel with which this state is, possibly, coupled strongly, and the threshold of which is situated below 600 MeV. In this energy region, only one channel is opened: this is the 4π channel. It is interesting to verify this assumption, because it concerns such an important state.

A minimum scenario of the simultaneous description of the processes $\pi\pi \rightarrow \pi\pi, K\bar{K}$ does not require the $f_0(1370)$ resonance; therefore, if this meson exists, it must be weakly coupled with the $\pi\pi$ channel, *e.g.* be the $s\bar{s}$ state (as to that assignment of the $f_0(1370)$ resonance, we agree with the work [24]).

The $f_0(1500)$ state is represented by the pole cluster which corresponds to a flavour singlet, *e.g.* the glueball.

We emphasize that the obtained results are model-independent, since they are based

on the first principles and on the mathematical fact that a local behaviour of analytic functions, determined on the Riemann surface, is governed by the nearest singularities on all sheets.

Finally, note that in the model-independent approach, there are many adjusted parameters (although, *e.g.* for the $\pi\pi$ scattering, they all are positions of poles describing resonances). The number of these parameters can be diminished by some dynamic assumptions, but this is another approach and of other value.

Acknowledgments

The authors are grateful to S. Dubníčka, S.B. Gerasimov, V.A. Meshcheryakov, V.N. Pervushin, M.K. Volkov and V.L. Yudichev for useful discussions and interest in this work.

This work has been supported by the Grant Program of Plenipotentiary of Slovak Republic at JINR. Yu.S. and M.N. were supported in part by the Slovak Scientific Grant Agency, Grant VEGA No. 2/7175/20; and D.K., by Grant VEGA No. 2/5085/99.

References

- [1] D. Krupa, V.A. Meshcheryakov, Yu.S. Surovtsev, *Nuovo Cimento A* 109 (1996) 281.
- [2] Review of Particle Physics, *Europ. Phys. J. C* 3 (1998) 1.
- [3] Y. Nambu, G. Jona-Lasinio, *Phys. Rev.* 122 (1961) 345.
- [4] M.K. Volkov, *Ann. Phys.* 157 (1984) 282.
- [5] T. Hatsuda, T. Kunihiro, *Phys. Rep.* 247 (1994) 223.
- [6] R. Delbourgo, M.D. Scadron, *Mod. Phys. Lett. A* 10 (1995) 251.
- [7] B.S. Zou, D.V. Bugg, *Phys. Rev. D* 48 (1993) R3948; *ibid.* D 50 (1994) 591.
- [8] M. Svec, *Phys. Rev. D* 53 (1996) 2343.
- [9] S. Ishida et al., *Progr. Theor. Phys.* 95 (1996) 745; *ibid.* 98 (1997) 621.
- [10] N. Törnqvist, *Phys. Rev. Lett.* 76 (1996) 1575.
- [11] R. Kamiński, L. Leśniak, B. Loiseau, *Eur. Phys. J. C* 9 (1999) 141.
- [12] D. Krupa, V.A. Meshcheryakov, Yu.S. Surovtsev, *Yad. Fiz.* 43 (1986) 231.; *Czech. J. Phys. B* 38 (1988) 1129.

- [13] D. Morgan, M.R. Pennington, Phys. Rev. D 48 (1993) 1185.
- [14] K.J. Le Couteur, Proc. Roy. Soc. A 256 (1960) 115.
- [15] R.G. Newton, J. Math. Phys. 2 (1961) 188.
- [16] M. Kato, Ann. Phys. **31**, 130 (1965).
- [17] J. Bohacik, H. Kühnelt, Phys. Rev. D 21 (1980) 1342.
- [18] B. Hyams et al., Nucl. Phys. B 64 (1973) 134; ibid. B 100 (1975) 205. A. Zylbersztejn et al., Phys. Lett. B 38 (1972) 457. P. Sonderegger, P. Bonamy, in: Proc. 5th Intern. Conf. on Elementary Particles, Lund, 1969, paper 372. J.R. Bensinger et al., Phys. Lett. B 36 (1971) 134. J.P. Baton et al., Phys. Lett. B 33 (1970) 525, 528. P. Baillon et al., Phys. Lett. B 38 (1972) 555. L. Rosselet et al., Phys. Rev. D 15 (1977) 574. A.A. Kartamyshev et al., Pis'ma v Zh. Eksp. Teor. Fiz. 25 (1977) 68. A.A. Bel'kov et al., Pis'ma v Zh. Eksp. Teor. Fiz. 29 (1979) 652.
- [19] A.B. Wicklund et al., Phys. Rev. Lett. 45 (1980) 1469. D. Cohen et al., Phys. Rev. D 22 (1980) 2595. A. Etkin et al., Phys. Rev. D 25 (1982) 1786.
- [20] S. Weinberg, Phys. Rev. Lett. 17 (1966) 616. B.W. Lee, H.T. Nieh, Phys. Rev. 166 (1968) 1507.
- [21] J. Gasser, H. Leutwyler, Ann. Phys. 158 (1984) 142.
- [22] M.K. Volkov, in: Physics of Elementary Particles and Atomic Nuclei, v.17, part 3 (1986) p. 433.
- [23] M.K. Volkov, V.L. Yudichev, M. Nagy, Nuovo Cimento A 112 (1999) 225.
- [24] L.S. Celenza, Huangsheng Wang, C.M. Shakin, Preprint of Brooklyn College of the City Univ. of New York, BCCNT: 00/041/289 (2000).

Figure Captions

Fig.1: Uniformization plane for the $\pi\pi$ -scattering amplitude.

Fig.2: The energy dependence of the ($|S_{12}|$) obtained on the basis of a simultaneous analysis of the experimental data on the coupled processes $\pi\pi \rightarrow \pi\pi, K\bar{K}$ in the channel with $I^G J^{PC} = 0^+ 0^{++}$. The data on the process $\pi\pi \rightarrow K\bar{K}$ are taken from Ref.[19].

Fig.3: The same as in Fig.2 but for the phase shift (δ_{12}).

Table 1: Pole clusters for obtained resonances

Sheet	$f_0(665)$		$f_0(980)$		$f_0(1500)$	
	E, MeV	Γ , MeV	E, MeV	Γ , MeV	E, MeV	Γ , MeV
II	610 ± 14	620 ± 26	988 ± 5	27 ± 8	1530 ± 25	390 ± 30
III	720 ± 15	55 ± 9	984 ± 16	210 ± 22	1430 ± 35	200 ± 30
					1510 ± 22	400 ± 34
IV					1410 ± 24	210 ± 38




Figure 1:




Figure 2:




Figure 3: